Leader in Biomolecular Solutions for Life Science

# **PPAT Rabbit pAb**

Catalog No.: A6698



## **Basic Information**

### **Observed MW**

57kDa

### **Calculated MW**

57kDa

### Category

Mouse Monoclonal Antibody

### **Applications**

WB,IF/ICC,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

The protein encoded by this gene is a member of the purine/pyrimidine phosphoribosyltransferase family. It is a regulatory allosteric enzyme that catalyzes the first step of de novo purine nucleotide biosythetic pathway. This gene and PAICS/AIRC gene, a bifunctional enzyme catalyzing steps six and seven of this pathway, are located in close proximity on chromosome 4, and divergently transcribed from an intergenic region.

# **Recommended Dilutions**

**WB** 1:500 - 1:2000

**IF/ICC** 1:50 - 1:100

# **Immunogen Information**

**Gene ID**5471

Swiss Prot
Q06203

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human PPAT (NP\_002694.3).

### **Synonyms**

GPAT; PRAT; ATASE; PPAT

### **Contact**

www.abclonal.com


# **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

# Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.




Western blot analysis of Rat spleen, using PPAT Rabbit pAb (A6698) at 1:2000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.



Immunofluorescence analysis of HeLa cells using PPAT Polyclonal Antibody (A6698) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.