Leader in Biomolecular Solutions for Life Science

# [KO Validated] TAGLN2 Rabbit pAb

ABclonal www.abclonal.com

Catalog No.: A6761

KO Validated

### **Basic Information**

# Observed MW

22kDa

#### **Calculated MW**

22kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB,IF/ICC,ELISA

#### **Cross-Reactivity**

Human, Rat

# **Background**

The protein encoded by this gene is similar to the protein transgelin, which is one of the earliest markers of differentiated smooth muscle. The specific function of this protein has not yet been determined, although it is thought to be a tumor suppressor. Multiple transcript variants encoding different isoforms have been found for this gene.

# **Recommended Dilutions**

WB

1:500 - 1:2000

IF/ICC

1:50 - 1:200

# **Immunogen Information**

**Gene ID** 

Swiss Prot P37802

8407

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-199 of human TAGLN2 (NP\_003555.1).

### **Synonyms**

HA1756; N2

#### **Contact**



www.abclonal.com

### **Product Information**

Source

Isotype

Purification

Rabbit

IgG


Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of lysates from wild type (WT) and TAGLN2 knockout (KO) 293T cells, using [KO Validated] TAGLN2 Rabbit pAb (A6761) at 1:1000 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5s.



Immunofluorescence analysis of C6 cells using [KO Validated] TAGLN2 Rabbit pAb (A6761) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.