Leader in Biomolecular Solutions for Life Science

# ZNF148 Rabbit pAb

Catalog No.: A7001 1 Publications



# **Basic Information**

## **Observed MW**

105kDa

### **Calculated MW**

89kDa

### Category

Polyclonal Antibody

### **Applications**

WB,IHC-P,IF/ICC,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

The protein encoded by this gene is a member of the Kruppel family of zinc finger DNA binding proteins. The encoded protein activates transcription of the T-cell receptor and intestinal alkaline phosphatase genes but represses transcription of the ornithine decarboxylase, vimentin, gastrin, stomelysin, and enolase genes. Increased expression of this gene results in decreased patient survival rates from colorectal cancer, while mutations in this gene have been associated with global developmental delay, hypoplastic corpus callosum, and dysmorphic facies.

# **Recommended Dilutions**

WB	1:500 - 1:1000
IHC-P	1:50 - 1:100
IF/ICC	1:20 - 1:100

# **Immunogen Information**

Gene ID	Swiss Prot
7707	Q9UQR1

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 690-794 of human ZNF148 (NP\_068799.2).

## **Synonyms**

BERF-1; BFCOL1; GDACCF; ZBP-89; ZFP148; pHZ-52; HT-BETA; ZNF148

## **Contact**

www.abclonal.com


# **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.




Western blot analysis of extracts of various cell lines, using ZNF148 Antibody (A7001) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 60s.




Immunohistochemistry analysis of paraffinembedded rat brain using ZNF148 antibody (A7001) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffinembedded human colon using ZNF148 antibody (A7001) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffinembedded mouse brain using ZNF148 antibody (A7001) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunofluorescence analysis of A-549 cells using ZNF148 antibody (A7001).