

A7134

Leader in Biomolecular Solutions for Life Science



## WDR77 Rabbit pAb

Catalog No.: A7134

### Basic Information

**Observed MW**

43kDa

**Calculated MW**

37kDa

**Category**

Polyclonal Antibody

**Applications**

WB,IF/ICC,ELISA

**Cross-Reactivity**

Human,Mouse,Rat

### Background

The protein encoded by this gene is an androgen receptor coactivator that forms a complex with protein arginine methyltransferase 5, which modifies specific arginines to dimethylarginines in several spliceosomal Sm proteins. The encoded protein may be involved in the early stages of prostate cancer, with most of the protein being nuclear-localized in benign cells but cytoplasmic in cancer cells. Several transcript variants encoding different isoforms have been found for this gene.

### Recommended Dilutions

**WB** 1:500 - 1:1000

**IF/ICC** 1:50 - 1:100

### Immunogen Information

**Gene ID**

79084

**Swiss Prot**

Q9BQA1

**Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 153-243 of human WDR77 (NP\_077007.1).

**Synonyms**

p44; MEP50; MEP-50; HKMT1069; Nbla10071; p44/Mep50; WDR77

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Product Information

**Source**

Rabbit

**Isotype**

IgG

**Purification**


Affinity purification

**Storage**

Store at -20°C. Avoid freeze / thaw cycles.



Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

Validation Data



Western blot analysis of extracts of various cell lines, using WDR77 antibody (A7134) at 1:1000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 1s.

Western blot analysis of extracts of Mouse brain, using WDR77 antibody (A7134) at 1:1000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 10s.



Immunofluorescence analysis of A-549 cells using WDR77 antibody (A7134). Blue: DAPI for nuclear staining.