Leader in Biomolecular Solutions for Life Science

# PLA2G2D Rabbit pAb

Catalog No.: A7165



### **Basic Information**

**Observed MW** 22kDa/

**Calculated MW** 17kDa

Category Polyclonal Antibody

Applications WB,IHC-P,IF/ICC,ELISA

**Cross-Reactivity** Human,Rat

#### Background

This gene encodes a secreted member of the phospholipase A2 family, and is found in a cluster of related family members on chromosome 1. Phospholipase A2 family members hydrolyze the sn-2 fatty acid ester bond of glycerophospholipids to produce lysophospholipids and free fatty acid. This gene may be involved in inflammation and immune response, and in weight loss associated with chronic obstructive pulmonary disease. Alternative splicing results in multiple transcript variants encoding different isoforms.

### **Recommended Dilutions**

### **Immunogen Information**

WB	1:500 - 1:2000	<b>Gene ID</b> 26279	Swiss Prot O9UNK4
IHC-P	1:50 - 1:200		Quality
IF/ICC	1:10 - 1:100	Immunogen Recombinant fusion protein containing a sequence corresponding to amino acids	

s 21-145 of human PLA2G2D (NP\_036532.1).

#### Synonyms

SPLASH; sPLA2S; PLA2IID; sPLA2-IID; PLA2G2D

### **Product Information**

€ www.abclonal.com

Isotype lgG

Purification Affinity purification


#### Storage

Source


Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

## **Validation Data**



Western blot analysis of various lysates using PLA2G2D Rabbit pAb (A7165) at 1:1000 dilution. Secondary antibody:HRP Goat Anti-Rabbit IgG (H+L)(AS014) at 1:10000 dilution. Lysates / proteins: 25 µg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection:ECL Basic Kit (RM00020). Exposuretime: 90s.



Immunofluorescence analysis of U2OS cells using PLA2G2D Rabbit pAb (A7165). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.