

A7257

Leader in Biomolecular Solutions for Life Science



Acetyl-Histone H3-K18 Rabbit pAb

Catalog No.: A7257

11 Publications

Basic Information

Observed MW

17kDa

Calculated MW

15kDa

Category

Mouse Monoclonal Antibody

Applications

WB, IHC-P, IF/ICC, ChIP, ELISA, DB

Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the small histone gene cluster on chromosome 6p22-p21.3.

Recommended Dilutions

WB	1:500 - 1:1000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200
ChIP	5µg antibody for 5µg-10µg of Chromatin
DB	1:500 - 1:1000

Immunogen Information

Gene ID
8290/8350

Swiss Prot
Q16695/P68431

Immunogen

A synthetic acetylated peptide around K18 of human H3 (NP_003520.1).

Synonyms

H3/j; H3C1; H3C2; H3C3; H3C4; H3C6; H3C7; H3C8; H3FJ; H3C10; H3C11; HIST1H3J; Acetyl-Histone H3-K18

Contact



www.abclonal.com

Product Information

Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.


Validation Data




Western blot analysis of lysates from HeLa cells using Acetyl-Histone H3-K18 Rabbit pAb (A7257) at 1:1000 dilution. HeLa cells were treated by TSA (1 μ M) at 37°C for 18 hours. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 0.5s.




Dot-blot analysis of all sorts of methylation peptides using Acetyl-Histone H3-K18 antibody (A7257) at 1:1000 dilution.




Immunohistochemistry analysis of Acetyl-Histone H3-K18 in paraffin-embedded rat brain using Acetyl-Histone H3-K18 Rabbit pAb (A7257) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.




Immunohistochemistry analysis of Acetyl-Histone H3-K18 in paraffin-embedded human gastric cancer using Acetyl-Histone H3-K18 Rabbit pAb (A7257) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of Acetyl-Histone H3-K18 in paraffin-embedded mouse brain using Acetyl-Histone H3-K18 Rabbit pAb (A7257) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.




Immunofluorescence analysis of C6 cells using Acetyl-Histone H3-K18 Rabbit pAb (A7257) at dilution of 1:100. C6 cells were treated by TSA (1 μ M) at 37°C for 18 hours. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of HeLa cells using Acetyl-Histone H3-K18 Rabbit pAb (A7257) at dilution of 1:100. HeLa cells were treated by TSA (1 μ M) at 37°C for 18 hours. Blue: DAPI for nuclear staining.

Validation Data



Immunofluorescence analysis of NIH/3T3 cells using Acetyl-Histone H3-K18 Rabbit pAb (A7257) at dilution of 1:100.NIH/3T3 cells were treated by TSA (1 uM) at 37°C for 18 hours. Blue: DAPI for nuclear staining.



Chromatin immunoprecipitation analysis of extracts of HCT116 cells, using Acetyl-Histone H3-K18 Rabbit pAb antibody (A7257) and rabbit IgG.The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.

