Leader in Biomolecular Solutions for Life Science

# **VLDLR Rabbit pAb**

Catalog No.: A7345 1 Publications



### **Basic Information**

### **Observed MW**

110kDa

### **Calculated MW**

96kDa

#### Category

Polyclonal Antibody

### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human, Mouse

### **Background**

The low density lipoprotein receptor (LDLR) gene family consists of cell surface proteins involved in receptor-mediated endocytosis of specific ligands. This gene encodes a lipoprotein receptor that is a member of the LDLR family and plays important roles in VLDL-triglyceride metabolism and the reelin signaling pathway. Mutations in this gene cause VLDLR-associated cerebellar hypoplasia. Alternative splicing generates multiple transcript variants encoding distinct isoforms for this gene.

### **Recommended Dilutions**

WB

1:500 - 1:2000

### **Immunogen Information**

Gene ID 7436 **Swiss Prot** 

P98155

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 833-869 of human VLDLR (NP\_003374.3).

### **Synonyms**

CAMRQ1; CARMQ1; CHRMQ1; VLDL-R; VLDLRCH; VLDLR

### **Contact**

•

www.abclonal.com

### **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of extracts of various cell lines, using VLDLR antibody (A7345) at 1:1000 dilution.\_Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.\_Lysates/proteins: 25 $\mu$ g per lane.\_Blocking buffer: 3% nonfat dry milk in TBST.\_Detection: ECL Enhanced Kit (RM00021).\_Exposure time: 90s.