

SYT4 Rabbit pAb

Catalog No.: A7737

Basic Information

Observed MW

46kDa

Calculated MW

48kDa

Category

Primary antibody

Applications

ELISA, WB, IHC-P, IF/ICC

Cross-Reactivity

Human, Mouse, Rat

Background

Predicted to enable several functions, including calcium ion binding activity; phospholipid binding activity; and syntaxin binding activity. Involved in negative regulation of catecholamine secretion and positive regulation of dendrite extension. Predicted to be located in several cellular components, including microvesicle; perinuclear region of cytoplasm; and secretory vesicle. Predicted to be active in several cellular components, including axon; exocytic vesicle; and glutamatergic synapse. Predicted to be integral component of neuronal dense core vesicle membrane.

Recommended Dilutions

WB	1:500 - 1:1000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200

Immunogen Information

Gene ID

6860

Swiss Prot

Q9H2B2

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 38-190 of human SYT4 (NP_065834.1).

Synonyms

HsT1192; SYT4

Contact

 | www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.


Validation Data




Western blot analysis of various lysates, using SYT4 Rabbit pAb (A7737) at 1:1000 dilution.
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 180s.



Immunofluorescence analysis of U2OS cells using SYT4 Rabbit pAb (A7737) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of SYT4 in paraffin-embedded human stomach using SYT4 Rabbit pAb (A7737) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of SYT4 in paraffin-embedded human liver cancer using SYT4 Rabbit pAb (A7737) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.