Leader in Biomolecular Solutions for Life Science

# [KO Validated] DDIT4 Rabbit pAb



Catalog No.: A8086

KO Validated

2 Publications

## **Basic Information**

## **Observed MW**

32kDa

### **Calculated MW**

25kDa

#### Category

Polyclonal Antibody

### **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human

# **Background**

Predicted to enable 14-3-3 protein binding activity. Involved in defense response to virus; negative regulation of TOR signaling; and response to hypoxia. Located in cytosol.

# **Recommended Dilutions**

WB

1:500 - 1:1000

# **Immunogen Information**

**Gene ID** 54541

Swiss Prot Q9NX09

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-232 of human DDIT4 ( $NP_061931.1$ ).

### **Synonyms**

Dig2; REDD1; REDD-1; T4

# **Contact**

•

www.abclonal.com

# **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of extracts from normal (control) and DDIT4 knockout (KO) HeLa cells, using DDIT4 antibody (A8086) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins:  $25\mu g$  per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 90s.

Western blot analysis of extracts of various cell lines, using DDIT4 antibody (A8086) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates proteins: 25µg per lane.

Biocking buffer: 3% nonfat dry milk in TBST.

7Detection: ECL Enhanced Kit (RM00021).

