Leader in Biomolecular Solutions for Life Science

CST6 Rabbit pAb

Catalog No.: A8115



Basic Information

Observed MW

15kDa

Calculated MW

17kDa

Category

Polyclonal Antibody

Applications

WB, IF/ICC, ELISA

Cross-Reactivity

Human

Background

The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins and the kininogens. The type 2 cystatin proteins are a class of cysteine proteinase inhibitors found in a variety of human fluids and secretions, where they appear to provide protective functions. This gene encodes a cystatin from the type 2 family, which is down-regulated in metastatic breast tumor cells as compared to primary tumor cells. Loss of expression is likely associated with the progression of a primary tumor to a metastatic phenotype.

Recommended Dilutions

WB 1:500 - 1:2000

IF/ICC 1:50 - 1:200

Immunogen Information

Gene IDSwiss Prot
1474
015828

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 29-149 of human CST6 (NP 001314.1).

Synonyms

ECTD15; CST6

Contact

www.abclonal.com


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification


Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of extracts of various cell lines, using CST6 antibody (A8115) at 1:1000 dilution._Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution._Lysates/proteins: 25 μ g per lane._Blocking buffer: 3% nonfat dry milk in TBST._Detection: ECL Enhanced Kit (RM00021)._Exposure time: 90s.



Immunofluorescence analysis of U2OS cells using CST6 antibody (A8115) at dilution of 1:100. Blue: DAPI for nuclear staining.