# ABclonal www.abclonal.com

## **CLDN8 Rabbit pAb**

Catalog No.: A8174 2 Publications

## **Basic Information**

#### **Observed MW**

28kDa

#### **Calculated MW**

25kDa

#### Category

Primary antibody

## **Applications**

ELISA,WB

#### **Cross-Reactivity**

Mouse, Rat

## **Background**

This gene encodes a member of the claudin family. Claudins are integral membrane proteins and components of tight junction strands. Tight junction strands serve as a physical barrier to prevent solutes and water from passing freely through the paracellular space between epithelial or endothelial cell sheets, and also play critical roles in maintaining cell polarity and signal transductions. This protein plays important roles in the paracellular cation barrier of the distal renal tubule, and in the paracellular barrier to prevent sodium back-leakage in distal colon. Differential expression of this gene has been observed in colorectal carcinoma and renal cell tumors, and along with claudin-7, is an immunohistochemical marker for the differential diagnosis of chromophobe renal cell carcinoma and renal oncocytoma.

## **Recommended Dilutions**

WB

1:500 - 1:2000

## **Immunogen Information**

Gene ID

**Swiss Prot** 

9073

P56748

## **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 29-81 of human CLDN8 (NP\_955360.1).

### **Synonyms**

HEL-S-79; CLDN8

#### **Contact**

•

www.abclonal.com

## **Product Information**

**Source** Rabbit Isotype

IgG


**Purification** 

Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of extracts of various cell lines, using CLDN8 antibody (A8174) at 1:3000

dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.