Leader in Biomolecular Solutions for Life Science

# **PDGFD Rabbit pAb**

Catalog No.: A8263



### **Basic Information**

### **Observed MW**

43kDa

### **Calculated MW**

43kDa

### Category

Polyclonal Antibody

### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human, Mouse

# **Background**

The protein encoded by this gene is a member of the platelet-derived growth factor family. The four members of this family are mitogenic factors for cells of mesenchymal origin and are characterized by a core motif of eight cysteines, seven of which are found in this factor. This gene product only forms homodimers and, therefore, does not dimerize with the other three family members. It differs from alpha and beta members of this family in having an unusual N-terminal domain, the CUB domain. Two splice variants have been identified for this gene.

# **Recommended Dilutions**

WB

1:500 - 1:2000

# **Immunogen Information**

**Gene ID** 80310

Swiss Prot Q9GZP0

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 250-350 of human PDGFD (NP 079484.1).

# **Synonyms**

IEGF; SCDGFB; MSTP036; SCDGF-B; PDGFD

### **Contact**

•

www.abclonal.com

### **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of extracts of various cell lines, using PDGFD antibody (A8263) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 90s.