

A8320

Leader in Biomolecular Solutions for Life Science



## NR1H4 Rabbit pAb

Catalog No.: A8320

11 Publications

### Basic Information

#### Observed MW

59kDa

#### Calculated MW

56kDa

#### Category

Polyclonal Antibody

#### Applications

WB, ELISA

#### Cross-Reactivity

Human, Mouse

### Background

This gene encodes a ligand-activated transcription factor that shares structural features in common with nuclear hormone receptor family members. This protein functions as a receptor for bile acids, and when bound to bile acids, binds to DNA and regulates the expression of genes involved in bile acid synthesis and transport. Alternatively spliced transcript variants encoding different isoforms have been described.

### Recommended Dilutions

WB 1:500 - 1:1000

### Immunogen Information

#### Gene ID

9971

#### Swiss Prot

Q96R11

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 207-476 of human NR1H4 (NP\_001193922.1).

#### Synonyms

BAR; FXR; HRR1; HRR-1; PFIC5; RIP14; NR1H4

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Product Information

#### Source

Rabbit

#### Isotype

IgG

#### Purification


Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.

## Validation Data



Western blot analysis of various lysates using NR1H4 Rabbit pAb (A8320) at 1:1000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (A5014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 10s.

Western blot analysis of lysates from RAW264.7 cells, using NR1H4 Rabbit pAb (A8320) at 1:1000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (A5014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 60s.

