## A8406

Leader in Biomolecular Solutions for Life Science

1 Publications

# COX7A2 Rabbit pAb

Catalog No.: A8406

**Basic Information** 

**Observed MW** 

**Calculated MW** 

Polyclonal Antibody

**Cross-Reactivity** 

Applications

11kDa

9kDa

Category

WB, ELISA

Human



# Background

Cytochrome c oxidase, the terminal component of the mitochondrial respiratory chain, catalyzes the electron transfer from reduced cytochrome c to oxygen. This component is a heteromeric complex consisting of three catalytic subunits encoded by mitochondrial genes, and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, while the nuclear-encoded subunits may function in the regulation and assembly of the complex. This nuclear gene encodes polypeptide 2 (liver isoform) of subunit VIIa, with this polypeptide being present in both muscle and non-muscle tissues. In addition to polypeptide 2, subunit VIIa includes polypeptide 1 (muscle isoform), which is present only in muscle tissues, and a related protein, which is present in all tissues. Alternative splicing results in multiple transcript variants. Related pseudogenes have been identified on chromosomes 4 and 14.

# **Recommended Dilutions**

## **Immunogen Information**

WB

# 1:500 - 1:2000

**Gene ID** 1347 Swiss Prot P14406

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 33-115 of human COX7A2 (NP 001856.2).

#### Synonyms

VIIAL; COX7AL; COX7AL1; COXVIIAL; COXVIIa-L; COX7A2

| Contact |
|---------|
|---------|

# **Product Information**

www.abclonal.com

**Isotype** IgG Purification Affinity purification


#### Storage

Source

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of extracts of HeLa cells, using COX7A2 antibody (A8406) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 10s.