Leader in Biomolecular Solutions for Life Science

IER3 Rabbit pAb

Catalog No.: A8566 1 Publications



Basic Information

Observed MW

17kDa

Calculated MW

17kDa

Category

Polyclonal Antibody

Applications

WB,IHC-P,ELISA

Cross-Reactivity

Human, Mouse

Background

This gene functions in the protection of cells from Fas- or tumor necrosis factor type alphainduced apoptosis. Partially degraded and unspliced transcripts are found after virus infection in vitro, but these transcripts are not found in vivo and do not generate a valid protein.

Recommended Dilutions

WB 1:100 - 1:500

IHC-P 1:50 - 1:100

Immunogen Information

Gene ID8870

Swiss Prot
P46695

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 50-150 of human IER3 (NP 003888.2).

Synonyms

DIF2; IEX1; PRG1; DIF-2; GLY96; IEX-1; IEX-1L; IER3

Contact

www.abclonal.com


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.




Western blot analysis of extracts of various cell lines, using IER3 antibody (A8566) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 180s.



Immunohistochemistry analysis of paraffinembedded mouse stomach using IER3 antibody (A8566) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffinembedded human gastric cancer using IER3 antibody (A8566) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.