Leader in Biomolecular Solutions for Life Science

# Periostin Rabbit mAb

Catalog No.: A9009 Recombinant 1 Publications



### **Basic Information**

### **Observed MW**

80kDa

### **Calculated MW**

80kDa/83kDa/84kDa/87kDa/90kDa/93kD

#### Category

SMab Recombinant Monoclonal Antibody

#### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human

### CloneNo number

ARC1380

# **Background**

This gene encodes a secreted extracellular matrix protein that functions in tissue development and regeneration, including wound healing, and ventricular remodeling following myocardial infarction. The encoded protein binds to integrins to support adhesion and migration of epithelial cells. This protein plays a role in cancer stem cell maintenance and metastasis. Mice lacking this gene exhibit cardiac valve disease, and skeletal and dental defects. Alternative splicing results in multiple transcript variants encoding different isoforms.

## **Recommended Dilutions**

WB

1:500 - 1:1000

## **Immunogen Information**

**Gene ID** 10631

Swiss Prot Q15063

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human Periostin (NP\_001129406.1).

### **Synonyms**

PN; OSF2; OSF-2; PDLPOSTN; Periostin

### **Contact**

•

www.abclonal.com

### **Product Information**

**Source** Rabbit **Purification**Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

Isotype

IgG

# **Validation Data**



Western blot analysis of extracts of MCF7 cells, using Periostin antibody (A9009) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.