A9909

Leader in Biomolecular Solutions for Life Science

3 Publications

PCNA Mouse mAb

Catalog No.: A9909



Background

The protein encoded by this gene is found in the nucleus and is a cofactor of DNA polymerase delta. The encoded protein acts as a homotrimer and helps increase the processivity of leading strand synthesis during DNA replication. In response to DNA damage, this protein is ubiquitinated and is involved in the RAD6-dependent DNA repair pathway. Two transcript variants encoding the same protein have been found for this gene. Pseudogenes of this gene have been described on chromosome 4 and on the X chromosome.

Basic Information

Observed MW 28kDa

Calculated MW 29kDa

Category Mouse Monoclonal Antibody

Applications WB,IHC-P,ELISA

Cross-Reactivity Human,Mouse,Rat

CloneNo number AMC0246

Recommended Dilutions

Immunogen Information

1:500 - 1:2000	Gene ID	Swiss Prot
	5111	P12004
1:50 - 1:200		

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-261 of human PCNA (P12004).

Synonyms ATLD2; PCNA

Contact

WB

IHC-P

Product Information


 www.abclonal.com

Source Mouse **lsotype** lgG2b,kappa Purification Affinity purification


Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates using PCNA Mouse mAb (A9909) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Mouse IgG (H+L) (AS003) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 60s.



Immunohistochemistry analysis of PCNA in paraffin-embedded human colon using PCNA Mouse mAb (A9909) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.