Leader in Biomolecular Solutions for Life Science

# MT-ND3 Rabbit pAb

Catalog No.: A9940 1 Publications



## **Basic Information**

#### **Observed MW**

13kDa

#### **Calculated MW**

13kDa

#### Category

Polyclonal Antibody

### **Applications**

WB,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

Predicted to enable NADH dehydrogenase (ubiquinone) activity. Predicted to be involved in cellular response to glucocorticoid stimulus; mitochondrial electron transport, NADH to ubiquinone; and response to oxidative stress. Located in mitochondrion. Is expressed in several structures, including alimentary system; brain; genitourinary system; hemolymphoid system gland; and liver and biliary system. Human ortholog(s) of this gene implicated in Leber hereditary optic neuropathy; Leigh disease; and Parkinson's disease. Orthologous to human MT-ND3 (mitochondrially encoded NADH:ubiquinone oxidoreductase core subunit 3).

# **Recommended Dilutions**

WB

1:500 - 1:2000

# **Immunogen Information**

**Gene ID** 17718

Swiss Prot

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 1-100 of mouse MT-ND3 (NP 904335.1).

### **Synonyms**

ND3; MT-ND3

### **Contact**

•

www.abclonal.com

# **Product Information**

**Source** Rabbit **Isotype** IgG Purification


Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of extracts of various cell lines, using MT-ND3 Antibody (A9940) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.