# ABclonal www.abclonal.com

## FBXO4 Rabbit pAb

Catalog No.: A9968

## **Basic Information**

#### **Observed MW**

44kDa

#### **Calculated MW**

44kDa

## Category

Primary antibody

## **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human, Mouse

## **Background**

This gene encodes a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of the ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbxs class. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

## **Recommended Dilutions**

WB

1:1000 - 1:4000

## **Immunogen Information**

**Gene ID** 26272

**Swiss Prot** 

Q9UKT5

## **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 138-387 of human FBXO4 (NP\_036308.1).

#### **Synonyms**

FBX4; FBXO4

## **Contact**

•

www.abclonal.com


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of extracts of various cell lines, using FBXO4 antibody (A9968) at 1:1000

dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 15s.