

## GAPDH Monoclonal Antibody

<b>Catalog No.</b>	AC002	<b>Category</b>	Loading control Antibodies
<b>Applications</b>	WB, IP	<b>Observed MW</b>	36kDa
<b>Cross-reactivity</b>	Human, Mouse, Rat	<b>Calculated MW</b>	31kDa/36kDa

### Immunogen Information

<b>Immunogen</b>	Recombinant protein of human GAPDH
<b>Gene ID</b>	2597
<b>Swiss prot</b>	P04406
<b>Synonyms</b>	GAPDH; G3PD; GAPD; HEL-S-162eP; glyceraldehyde -3-phosphate dehydrogenase

### Product information

<b>Source</b>	Mouse
<b>Isotype</b>	IgG
<b>Purification method</b>	Affinity purification
<b>Storage</b>	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

### Background

This gene encodes a member of the glyceraldehyde-3-phosphate dehydrogenase protein family. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. The product of this gene catalyzes an important energy-yielding step in carbohydrate metabolism, the reversible oxidative phosphorylation of glyceraldehyde-3-phosphate in the presence of inorganic phosphate and nicotinamide adenine dinucleotide (NAD). The encoded protein has additionally been identified to have uracil DNA glycosylase activity in the nucleus. Also, this protein contains a peptide that has antimicrobial activity against *E. coli*, *P. aeruginosa*, and *C. albicans*. Studies of a similar protein in mouse have assigned a variety of additional functions including nitrosylation of nuclear proteins, the regulation of mRNA stability, and acting as a transferrin receptor on the cell surface of macrophage. Many pseudogenes similar to this locus are present in the human genome. Alternative splicing results in multiple transcript variants.

### Recommended Dilutions

WB 1:5000 -  
1:20000  
IP 1:50 - 1:200



Western blot - GAPDH Monoclonal Antibody (AC002)