

AE017

Leader in Biomolecular Solutions for Life Science



## Mouse anti V5-Tag mAb

Catalog No.: AE017

11 Publications

### Basic Information

#### Observed MW

Refer to Figures

#### Calculated MW

#### Category

Polyclonal Antibody

#### Applications

WB,IF/ICC,IP,ELISA

#### Cross-Reactivity

Species independent

#### CloneNo number

AMC0506

### Background

Protein tags are peptide sequences genetically grafted onto a recombinant protein. Often these tags are removable by chemical agents or by enzymatic means, such as proteolysis or intein splicing. Tags are attached to proteins for various purposes. Epitope tags are short peptide sequences which are chosen because high-affinity antibodies can be reliably produced in many different species. These are usually derived from viral genes, which explain their high immunoreactivity. Epitope tags include V5-tag, Myc-tag, HA-tag and NE-tag. These tags are particularly useful for western blotting, immunofluorescence and immunoprecipitation experiments, although they also find use in antibody purification.

### Recommended Dilutions

WB 1:1000 - 1:5000

IF/ICC 1:50 - 1:200

IP 0.5µg-4µg antibody for  
200µg-400µg extracts of  
whole cells

### Immunogen Information

#### Gene ID

#### Swiss Prot

#### Immunogen

A synthetic peptide corresponding to V5 tag.

#### Synonyms

V5;V5 tag;V5-tag

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Product Information

#### Source

Mouse

#### Isotype

IgG2b, Kappa

#### Purification


Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.

## Validation Data



Western blot analysis of recombinant Mouse V5 protein using Mouse anti V5-Tag mAb (AE017) at dilution of 1:2000.  
 Secondary antibody: HRP Goat Anti-Mouse IgG (H+L) (AS003) at 1:10000 dilution.  
 Lysates/proteins: 10ng per lane.  
 Blocking buffer: 3% nonfat dry milk in TBST.  
 Detection: ECL Basic Kit (RM00020).  
 Exposure time: 1s.

Immunoprecipitation analysis of 300ug extracts of CHO cells using 3ug Mouse anti V5-Tag mAb antibody (AE017 1:50). Western blot was performed from the immunoprecipitate using Mouse anti V5-Tag mAb antibody (AE017) at a dilution of 1:2000.

