Leader in Biomolecular Solutions for Life Science

# Phospho-ErbB4/HER4-Y1284 Rabbit pAb



Catalog No.: AP0034

1 Publications

### **Basic Information**

#### **Observed MW**

180kDa

#### **Calculated MW**

147kDa

#### Category

Mouse Monoclonal Antibody

#### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

This gene is a member of the Tyr protein kinase family and the epidermal growth factor receptor subfamily. It encodes a single-pass type I membrane protein with multiple cysteine rich domains, a transmembrane domain, a tyrosine kinase domain, a phosphotidylinositol-3 kinase binding site and a PDZ domain binding motif. The protein binds to and is activated by neuregulins and other factors and induces a variety of cellular responses including mitogenesis and differentiation. Multiple proteolytic events allow for the release of a cytoplasmic fragment and an extracellular fragment. Mutations in this gene have been associated with cancer. Alternatively spliced variants which encode different protein isoforms have been described; however, not all variants have been fully characterized.

# **Recommended Dilutions**

WB

1:500 - 1:1000

### Immunogen Information

Gene ID

Swiss Prot 015303

# Immunogen

A synthetic phosphorylated peptide around Y1284 of human ErbB4/HER4 (NP\_005226.1).

#### Synonyms

HER4; ALS19; p180erbB4; Phospho-ErbB4/HER4-Y1284

### **Contact**

•

www.abclonal.com

### **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of lysates from MCF7 cells using Phospho-ErbB4/HER4-Y1284 Rabbit pAb (AP0034) at 1:700 dilution. MCF7 cells were treated by IGF (200 ng/ml) at 37°C for 30 minutes after serumstarvation overnight.

Secondary antibody:HRP Goat Anti-Rabbit IgG (H+L)(AS014) at 1:10000 dilution.

Lysates/proteins: 20 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection:ECL Basic Kit (RM00020).

Exposuretime: 30s.