

# Phospho-GSK3 $\beta$ -S9 Rabbit pAb

Catalog No.: AP0039 **29 Publications**

## Basic Information

### Observed MW

46 kDa

### Calculated MW

47kDa

### Category

Primary antibody

### Applications

ELISA, WB, IHC-P, IF/ICC

### Cross-Reactivity

Human, Mouse, Rat

## Background

The protein encoded by this gene is a serine-threonine kinase belonging to the glycogen synthase kinase subfamily. It is a negative regulator of glucose homeostasis and is involved in energy metabolism, inflammation, ER-stress, mitochondrial dysfunction, and apoptotic pathways. Defects in this gene have been associated with Parkinson disease and Alzheimer disease.

## Recommended Dilutions

<b>WB</b>	1:100 - 1:500
<b>IHC-P</b>	1:50 - 1:200
<b>IF/ICC</b>	1:50 - 1:200

## Immunogen Information

<b>Gene ID</b>	<b>Swiss Prot</b>
2932	P49841

### Immunogen

A synthetic phosphorylated peptide around S9 of human GSK3 $\beta$  (NP\_001139628.1).

### Synonyms

GSK3B; gsk-3 $\beta$ ; Phospho-GSK3 $\beta$ -S9

## Contact

 | [www.abclonal.com](http://www.abclonal.com)

## Product Information


<b>Source</b>	<b>Isotype</b>	<b>Purification</b>
Rabbit	IgG	Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.


Validation Data




Western blot analysis of lysates from C6 cells using Phospho-GSK3β-S9 Rabbit pAb (AP0039) at 1:500 dilution. C6 cells were treated by Calyculin A (100 nM) at 37°C for 30 minutes after serum-starvation overnight.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25 µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 10s.




Immunohistochemistry analysis of Phospho-GSK3β-S9 in paraffin-embedded human colon tissue using Phospho-GSK3β-S9 Rabbit pAb (AP0039) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of Phospho-GSK3β-S9 in paraffin-embedded human colon carcinoma tissue using Phospho-GSK3β-S9 Rabbit pAb (AP0039) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of Phospho-GSK3β-S9 in paraffin-embedded human liver cancer tissue using Phospho-GSK3β-S9 Rabbit pAb (AP0039) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunofluorescence analysis of HeLa cells using Phospho-GSK3β-S9 Rabbit pAb (AP0039) at a dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using Phospho-GSK3β-S9 Rabbit pAb (AP0039) at a dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of PC-12 cells using Phospho-GSK3β-S9 Rabbit pAb (AP0039) at a dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.