

AP0115

Leader in Biomolecular Solutions for Life Science



## Phospho-mTOR-S2448 Rabbit mAb

Catalog No.: AP0115

Recombinant

30 Publications

### Basic Information

#### Observed MW

289kDa

#### Calculated MW

289kDa

#### Category

SMab Recombinant Monoclonal Antibody

#### Applications

WB, ELISA

#### Cross-Reactivity

Human, Mouse, Rat

#### CloneNo number

ARC0094

#### Conjugate

Unconjugated

### Recommended Dilutions

WB 1:500 - 1:1000

### Background

The protein encoded by this gene belongs to a family of phosphatidylinositol kinase-related kinases. These kinases mediate cellular responses to stresses such as DNA damage and nutrient deprivation. This kinase is a component of two distinct complexes, mTORC1, which controls protein synthesis, cell growth and proliferation, and mTORC2, which is a regulator of the actin cytoskeleton, and promotes cell survival and cell cycle progression. This protein acts as the target for the cell-cycle arrest and immunosuppressive effects of the FKBP12-rapamycin complex. Inhibitors of mTOR are used in organ transplants as immunosuppressants, and are being evaluated for their therapeutic potential in SARS-CoV-2 infections. Mutations in this gene are associated with Smith-Kingsmore syndrome and somatic focal cortical dysplasia type II. The ANGPTL7 gene is located in an intron of this gene.

### Immunogen Information

#### Gene ID

2475

#### Swiss Prot

P42345

#### Immunogen

A synthetic phosphorylated peptide around S2448 of human mTOR (P42345).

#### Synonyms

SKS; FRAP; FRAP1; FRAP2; RAFT1; RAPT1; Phospho-mTOR-S2448

### Product Information

#### Source

Rabbit

#### Isotype

IgG

#### Purification

Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

### Contact




[www.abclonal.com](http://www.abclonal.com)

Validation Data



Western blot analysis of extracts of 293T cells, using Phospho-mTOR-S2448 Rabbit mAb (AP0115) at 1:1000 dilution. 293T cells were treated by 10% FBS at 37°C for 30 minutes after serum-starvation overnight or treated by IGF-1 (50 ng/mL) at 37°C for 5 minutes after serum-starvation overnight. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1s.



Western blot analysis of extracts of NIH/3T3 cells, using Phospho-mTOR-S2448 antibody (AP0115) at 1:1000 dilution. NIH/3T3 cells were treated by EGF (100 ng/ml) at 37°C for 30 minutes after serum-starvation overnight. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 90s.