Leader in Biomolecular Solutions for Life Science

# Phospho-FAK-Y397 Rabbit pAb

Catalog No.: AP0302 19 Publications



### **Basic Information**

### **Observed MW**

125kDa

### **Calculated MW**

119kDa

#### Category

Mouse Monoclonal Antibody

### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

This gene encodes a cytoplasmic protein tyrosine kinase which is found concentrated in the focal adhesions that form between cells growing in the presence of extracellular matrix constituents. The encoded protein is a member of the FAK subfamily of protein tyrosine kinases but lacks significant sequence similarity to kinases from other subfamilies. Activation of this gene may be an important early step in cell growth and intracellular signal transduction pathways triggered in response to certain neural peptides or to cell interactions with the extracellular matrix. Several transcript variants encoding different isoforms have been found for this gene.

# **Recommended Dilutions**

WB

1:100 - 1:500

## Immunogen Information

Gene ID

Swiss Prot 005397

#### **Immunogen**

A synthetic phosphorylated peptide around Y397 of human FAK (NP\_722560.1).

#### Synonyms

FAK; FADK; FAK1; FRNK; FADK 1; PPP1R71; p125FAK; pp125FAK; Phospho-FAK-Y397

### **Contact**

•

www.abclonal.com

### **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of lysates from A-549 cells, using Phospho-FAK-Y397 Rabbit pAb (AP0302) at 1:400

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.