Leader in Biomolecular Solutions for Life Science

# Phospho-VEGF Receptor 2-Y1175 Rabbit pAb



Catalog No.: AP0382

16 Publications

## **Basic Information**

# Observed MW

230kDa

#### **Calculated MW**

152kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

Vascular endothelial growth factor (VEGF) is a major growth factor for endothelial cells. This gene encodes one of the two receptors of the VEGF. This receptor, known as kinase insert domain receptor, is a type III receptor tyrosine kinase. It functions as the main mediator of VEGF-induced endothelial proliferation, survival, migration, tubular morphogenesis and sprouting. The signalling and trafficking of this receptor are regulated by multiple factors, including Rab GTPase, P2Y purine nucleotide receptor, integrin alphaVbeta3, T-cell protein tyrosine phosphatase, etc.. Mutations of this gene are implicated in infantile capillary hemangiomas.

# **Recommended Dilutions**

WB

1:100 - 1:500

## **Immunogen Information**

Gene ID

Swiss Prot P35968

#### **Immunogen**

A synthetic phosphorylated peptide around Y1175 of human VEGF Receptor 2 (NP 002244.1).

## **Synonyms**

FLK1; CD309; VEGFR; VEGFR2; Phospho-VEGF Receptor 2-Y1175

## **Contact**

•

www.abclonal.com

## **Product Information**

**Source** Rabbit **Isotype** IgG


**Purification**Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of A-431, using Phospho-VEGF Receptor 2-Y1175 antibody (AP0382) at 1:500 dilution.A-431 cells were treated by EGF (100 ng/ml) at  $37^{\circ}$ C for 30 minutes after serum-starvation overnight.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.