Leader in Biomolecular Solutions for Life Science

# Phospho-TP73-Y99 Rabbit pAb

Catalog No.: AP0465



### **Basic Information**

### **Observed MW**

80kDa

### **Calculated MW**

70kDa

### Category

Polyclonal Antibody

### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human

# **Background**

This gene encodes a member of the p53 family of transcription factors involved in cellular responses to stress and development. It maps to a region on chromosome 1p36 that is frequently deleted in neuroblastoma and other tumors, and thought to contain multiple tumor suppressor genes. The demonstration that this gene is monoallelically expressed (likely from the maternal allele), supports the notion that it is a candidate gene for neuroblastoma. Many transcript variants resulting from alternative splicing and/or use of alternate promoters have been found for this gene, but the biological validity and the full-length nature of some variants have not been determined.

# **Recommended Dilutions**

WB

1:500 - 1:2000

# **Immunogen Information**

Gene ID

Swiss Prot 015350

#### **Immunogen**

A phospho specific peptide corresponding to residues surrounding Y99 of human TP73

# **Synonyms**

P73; CILD47; Phospho-TP73-Y99

### **Contact**

•

www.abclonal.com

# **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

# Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of MCF7, using Phospho-TP73-Y99 Rabbit pAb (AP0465) at 1:500 dilution.MCF7 cells were treated by pervanadate (1 mM) at 37°C for 30 minutes after serum-starvation overnight.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 45s.