Leader in Biomolecular Solutions for Life Science

# Phospho-PDK1/PDPK1-S241 Rabbit pAb



Catalog No.: AP0477

1 Publications

### **Basic Information**

### **Observed MW**

60-70kDa

### **Calculated MW**

63kDa

### Category

Polyclonal Antibody

### **Applications**

WB,IF/ICC,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

Enables 3-phosphoinositide-dependent protein kinase activity; phospholipase activator activity; and phospholipase binding activity. Involved in several processes, including cell surface receptor signaling pathway; regulation of protein kinase activity; and regulation of signal transduction. Acts upstream of or within intracellular signal transduction. Located in cell projection; cytosol; and plasma membrane. Implicated in prostate cancer. Biomarker of lung non-small cell carcinoma.

# **Recommended Dilutions**

**WB** 1:500 - 1:2000

IF/ICC 1:50 - 1:200

# **Immunogen Information**

**Gene ID Swiss Prot** 5170 015530

#### **Immunogen**

A synthetic phosphorylated peptide around S241 of human PDK1/PDPK1 (NP\_002604.1).

#### Synonyms

PDK1; PDPK2; PDPK2P; PRO0461; Phospho-PDK1/PDPK1-S241

### **Contact**

www.abclonal.com

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# **Validation Data**




Western blot analysis of extracts of NIH/3T3 cells, using Phospho-PDK1/PDPK1-S241 pAb (AP0477) at 1:1000 dilution.NIH/3T3 cells were treated by PDGF (100 ng/mL) at  $37^{\circ}\text{C}$  for 30 minutes after serum-starvation overnight.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25µg per lane. Blocking buffer: 3% BSA.

Detection: ECL Basic Kit (RM00020).


Exposure time: 60s.



Immunofluorescence analysis of C6 cells using Phospho-PDK1/PDPK1-S241 Rabbit pAb (AP0477) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH-3T3 cells using Phospho-PDK1/PDPK1-S241 Rabbit pAb (AP0477) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U-2 OS cells using Phospho-PDK1/PDPK1-S241 Rabbit pAb (AP0477) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.