

AP0505

Leader in Biomolecular Solutions for Life Science



Phospho-IKK α -S176 Rabbit pAb

Catalog No.: AP0505

6 Publications

Basic Information

Observed MW

84kDa

Calculated MW

85kDa

Category

Mouse Monoclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a member of the serine/threonine protein kinase family. The encoded protein, a component of a cytokine-activated protein complex that is an inhibitor of the essential transcription factor NF-kappa-B complex, phosphorylates sites that trigger the degradation of the inhibitor via the ubiquitination pathway, thereby activating the transcription factor.

Recommended Dilutions

WB 1:500 - 1:2000

Immunogen Information

Gene ID

1147

Swiss Prot

O15111

Immunogen

A synthetic phosphorylated peptide around S176 of human IKK α (NP_001269.3).

Synonyms

BPS2; IKK1; IKKA; IKBKA; TCF16; NFKBIKA; IKK-alpha; Phospho-IKK α -S176

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH 7.3.

Validation Data



Western blot analysis of lysates from C6 cells, using Phospho-IKK α -S176 Rabbit pAb (AP0505) at 1:1000 dilution. C6 cells were treated by TNF- α (20 ng/mL) at 37°C for 30 minutes.
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25 μ g per lane.
 Blocking buffer: 3% BSA.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 30s.

Western blot analysis of lysates from HeLa cells, using Phospho-IKK α -S176 Rabbit pAb (A0423). HeLa cells were treated by TNF- α (20 ng/mL) at 37°C for 30 minutes.
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25 μ g per lane.
 Blocking buffer: 3% BSA.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 30s.

