

AP0546

Leader in Biomolecular Solutions for Life Science



Phospho-IKK α / β -S176/180 Rabbit pAb

Catalog No.: AP0546

8 Publications

Basic Information

Observed MW

85kDa

Calculated MW

84kDa/29kDa/79kDa/85kDa/86kDa

Category

Mouse Monoclonal Antibody

Applications

WB, IHC-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a member of the serine/threonine protein kinase family. The encoded protein, a component of a cytokine-activated protein complex that is an inhibitor of the essential transcription factor NF-kappa-B complex, phosphorylates sites that trigger the degradation of the inhibitor via the ubiquination pathway, thereby activating the transcription factor. [provided by RefSeq, Jul 2008]

Recommended Dilutions

WB	1:500 - 1:1000
IHC-P	1:100 - 1:500

Immunogen Information

Gene ID

1147/3551

Swiss Prot

O15111/O14920

Immunogen

A synthetic phosphorylated peptide around S176 & S180 of human IKK-alphaCHUK (NP_001269.3).

Synonyms

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of lysates from HeLa cells, using Phospho-IKKα/β-S176/180 Rabbit pAb (AP0546) at 1:1000 dilution. HeLa cells were treated by TNF-α (20 ng/ml) at 37°C for 30 minutes after serum-starvation overnight.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25μg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Enhanced Kit (RM00021).


Exposure time: 90s.




Immunohistochemistry analysis of Phospho-IKKα/β-S176/180 in paraffin-embedded Human colon carcinoma tissue using Phospho-IKKα/β-S176/180 Rabbit pAb (AP0546) at a dilution of 1:300 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of Phospho-IKKα/β-S176/180 in paraffin-embedded Human pancreas tissue using Phospho-IKKα/β-S176/180 Rabbit pAb (AP0546) at a dilution of 1:300 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of Phospho-IKKα/β-S176/180 in paraffin-embedded Mouse kidney tissue using Phospho-IKKα/β-S176/180 Rabbit pAb (AP0546) at a dilution of 1:300 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of Phospho-IKKα/β-S176/180 in paraffin-embedded Rat kidney tissue using Phospho-IKKα/β-S176/180 Rabbit pAb (AP0546) at a dilution of 1:300 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of Phospho-IKKα/β-S176/180 in paraffin-embedded Human lung cancer tissue using Phospho-IKKα/β-S176/180 Rabbit pAb (AP0546) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of Phospho-IKKα/β-S176/180 in paraffin-embedded Human pancreas tissue using Phospho-IKKα/β-S176/180 Rabbit pAb (AP0546) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.

Validation Data



Immunohistochemistry analysis of Phospho-IKKα/β-S176/180 in paraffin-embedded Mouse kidney tissue using Phospho-IKKα/β-S176/180 Rabbit pAb (AP0546) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of Phospho-IKKα/β-S176/180 in paraffin-embedded Rat kidney tissue using Phospho-IKKα/β-S176/180 Rabbit pAb (AP0546) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.