

AP0554

Leader in Biomolecular Solutions for Life Science



## Phospho-Smad3-T179 Rabbit pAb

Catalog No.: AP0554

5 Publications

### Basic Information

#### Observed MW

48kDa

#### Calculated MW

48kDa

#### Category

Polyclonal Antibody

#### Applications

WB, IHC-P, ELISA

#### Cross-Reactivity

Human, Mouse, Rat

### Background

The SMAD family of proteins are a group of intracellular signal transducer proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. The SMAD3 protein functions in the transforming growth factor-beta signaling pathway, and transmits signals from the cell surface to the nucleus, regulating gene activity and cell proliferation. This protein forms a complex with other SMAD proteins and binds DNA, functioning both as a transcription factor and tumor suppressor. Mutations in this gene are associated with aneurysms-osteoarthritis syndrome and Loews-Dietz Syndrome 3.

### Recommended Dilutions

WB 1:500 - 1:2000

IHC-P 1:50 - 1:200

### Immunogen Information

#### Gene ID

4088

#### Swiss Prot

P84022

#### Immunogen

A synthetic phosphorylated peptide around T179 of human Smad3 (NP\_005893.1).

#### Synonyms

LDS3; mad3; LDS1C; MADH3; JV15-2; hMAD-3; hSMAD3; HSPC193; HsT17436; Phospho-Smad3-T179

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Product Information

#### Source

Rabbit

#### Isotype

IgG

#### Purification


Affinity purification

#### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.


Validation Data




Western blot analysis of extracts of HeLa cell line, using Phospho-Smad3-T179 antibody (AP0554). The lane on the right is treated with the antigen-specific peptide.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% BSA.




Western blot analysis of extracts of PC-12 cells, using Phospho-Smad2-S465/467/-Smad3-S423/425 pAb (AP0554) at 1:1000 dilution. PC-12 cells were treated by TNF-α (20 ng/mL) at 37°C for 30 minutes after serum starvation overnight.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% BSA.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 30s.



Immunohistochemistry analysis of paraffin-embedded human breast using Phospho-Smad3-T179 Rabbit pAb (AP0554) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of paraffin-embedded human colon using Phospho-Smad3-T179 Rabbit pAb (AP0554) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse kidney using Phospho-Smad3-T179 Rabbit pAb (AP0554) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

## Validation Data

---



Immunohistochemistry analysis of paraffin-embedded rat liver using Phospho-Smad3-T179 Rabbit pAb (AP0554) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.