Leader in Biomolecular Solutions for Life Science

ABclonal www.abclonal.com

Phospho-PLC gamma 2 (PLCG2)-Y759 Rabbit pAb

Catalog No.: AP0785 1 Publications

Basic Information

Observed MW

150kDa

Calculated MW

148kDa

Category

Polyclonal Antibody

Applications

WB,ELISA

Cross-Reactivity

Human

Background

The protein encoded by this gene is a transmembrane signaling enzyme that catalyzes the conversion of 1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate to 1D-myo-inositol 1,4,5-trisphosphate (IP3) and diacylglycerol (DAG) using calcium as a cofactor. IP3 and DAG are second messenger molecules important for transmitting signals from growth factor receptors and immune system receptors across the cell membrane. Mutations in this gene have been found in autoinflammation, antibody deficiency, and immune dysregulation syndrome and familial cold autoinflammatory syndrome 3.

Recommended Dilutions

WB

1:500 - 1:2000

Immunogen Information

Gene ID 5336 Swiss Prot P16885

Immunogen

A synthetic phosphorylated peptide around Y759 of human PLC gamma 2 (PLCG2) (NP 002652.2).

Synonyms

FCAS3; APLAID; PLC-IV; PLC-gamma-2; Phospho-PLC gamma 2 (PLCG2)-Y759

Contact

•

www.abclonal.com

Product Information


Source Rabbit **Isotype** IgG **Purification**Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data



Western blot analysis of Raji, using Phospho-PLC gamma 2 (PLCG2)-Y759 Rabbit pAb (AP0785) at 1:2000 dilution.Raji cells were treated by Pervanadate (1 mM) at 37°C for 30 minutes.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.