AP0979

Leader in Biomolecular Solutions for Life Science

Phospho-β-Catenin-S33/S37 Rabbit mAb

Catalog No.: AP0979

Recombinant 2 Publications



Basic Information

Observed MW 100kDa

Calculated MW 85kDa

Category SMab Recombinant Monoclonal Antibody

Applications WB,ELISA

Cross-Reactivity Human

CloneNo number ARC1523

Background

The protein encoded by this gene is part of a complex of proteins that constitute adherens junctions (AJs). AJs are necessary for the creation and maintenance of epithelial cell layers by regulating cell growth and adhesion between cells. The encoded protein also anchors the actin cytoskeleton and may be responsible for transmitting the contact inhibition signal that causes cells to stop dividing once the epithelial sheet is complete. Finally, this protein binds to the product of the APC gene, which is mutated in adenomatous polyposis of the colon. Mutations in this gene are a cause of colorectal cancer (CRC), pilomatrixoma (PTR), medulloblastoma (MDB), and ovarian cancer. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

Immunogen Information

WB

1:500 - 1:1000

Gene ID 1499 Swiss Prot P35222

Immunogen

A synthetic phosphorylated peptide around S33 of human beta CateninCTNB1 (P35222).

Synonyms

EVR7; CTNNB; MRD19; NEDSDV; armadillo; Phospho-β-Catenin-S33/S37

Contact

Product Information

www.abclonal.com

Purification Affinity purification

Storage


Source

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.

Isotype

lgG



Western blot analysis of extracts of 293T cells, using Phospho-β-Catenin-S33/S37 Rabbit mAb (AP0979) at 1:1000 dilution.293T cells were treated by Calyculin A (100 nM) at 37°C for 30 minutes after serumstarvation overnight. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% BSA. Detection: ECL Basic Kit (RM00020). Exposure time: 3min.