

AP1026

Leader in Biomolecular Solutions for Life Science



Phospho-TBK1/NAK-S172 Rabbit mAb

Catalog No.: AP1026

Recombinant

17 Publications

Basic Information

Observed MW

84kDa

Calculated MW

84kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse

CloneNo number

ARC1571

Background

The NF-kappa-B (NFKB) complex of proteins is inhibited by I-kappa-B (IKB) proteins, which inactivate NFKB by trapping it in the cytoplasm. Phosphorylation of serine residues on the IKB proteins by IKB kinases marks them for destruction via the ubiquitination pathway, thereby allowing activation and nuclear translocation of the NFKB complex. The protein encoded by this gene is similar to IKB kinases and can mediate NFKB activation in response to certain growth factors. The protein is also an important kinase for antiviral innate immunity response.

Recommended Dilutions

WB 1:500 - 1:2000

Immunogen Information

Gene ID

29110

Swiss Prot

Q9UHD2

Immunogen

A synthetic phosphorylated peptide around S172 of human TBK1/NAKTBK1 (Q9UHD2).

Synonyms

NAK; T2K; IIAE8; FTDALS4; Phospho-TBK1/NAK-S172

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of various lysates using Phospho-TBK1/NAK-S172 Rabbit mAb (AP1026) at 1:1000 dilution. Both HeLa cells and NIH/3T3 cells were treated by Calyculin A (100 nM) at 37°C for 30 minutes after serum-starvation overnight.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 μ g per lane.
Blocking buffer: 3% BSA.
Detection: ECL Basic Kit (RM00020).
Exposure time: 1min.