

AP1119

Leader in Biomolecular Solutions for Life Science



Phospho-Vimentin-S83 Rabbit pAb

Catalog No.: AP1119

Basic Information

Observed MW

57kDa

Calculated MW

54kDa

Category

Polyclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a type III intermediate filament protein. Intermediate filaments, along with microtubules and actin microfilaments, make up the cytoskeleton. The encoded protein is responsible for maintaining cell shape and integrity of the cytoplasm, and stabilizing cytoskeletal interactions. This protein is involved in neuritogenesis and cholesterol transport and functions as an organizer of a number of other critical proteins involved in cell attachment, migration, and signaling. Bacterial and viral pathogens have been shown to attach to this protein on the host cell surface. Mutations in this gene are associated with congenital cataracts in human patients.

Recommended Dilutions

WB 1:500 - 1:1000

Immunogen Information

Gene ID

7431

Swiss Prot

P08670

Immunogen

A synthetic phosphorylated peptide around S83 of human VIM (NP_003371.2).

Synonyms

VIM; CTRCT30; HEL113; vimentin; Phospho-Vimentin-S83

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.


Validation Data



Western blot analysis of extracts of NIH/3T3 cells, using Phospho-Vimentin-S83 antibody (AP1119) at 1:1000 dilution. NIH/3T3 cells were treated by Paclitaxel (100 nM/ml) at 37°C for 20 hours. NIH/3T3 cells were treated by Hydroxyurea (4 mM) at 37°C for 20 hours. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1s.



Western blot analysis of extracts of C6 cells, using Phospho-Vimentin-S83 antibody (AP1119) at 1:1000 dilution. C6 cells were treated by Paclitaxel (100 nM) at 37°C for 20 hours. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.



Western blot analysis of extracts of HeLa cells, using Phospho-Vimentin-S83 antibody (AP1119) at 1:1000 dilution. HeLa cells were treated by Paclitaxel (100 nM/ml) at 37°C for 20 hours. HeLa cells were treated by Hydroxyurea (4 mM) at 37°C for 20 hours. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.