

# Phospho-ERK1-T202/Y204 + ERK2-T185/Y187 Rabbit pAb

www.abclonal.com

Catalog No.: AP1120 6 Publications

### **Basic Information**

### **Observed MW**

42kDa/44kDa

### **Calculated MW**

### Category

Primary antibody

### **Applications**

ELISA,WB

### **Cross-Reactivity**

Human, Mouse

# **Background**

This gene encodes a member of the MAP kinase family. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. The activation of this kinase requires its phosphorylation by upstream kinases. Upon activation, this kinase translocates to the nucleus of the stimulated cells, where it phosphorylates nuclear targets. One study also suggests that this protein acts as a transcriptional repressor independent of its kinase activity. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. Two alternatively spliced transcript variants encoding the same protein, but differing in the UTRs, have been reported for this gene. [provided by RefSeq, Jan 2014]

# **Recommended Dilutions**

WB

1:500 - 1:2000

# **Immunogen Information**

Gene ID

**Swiss Prot** 

5594/5595

P28482/P27361

### **Immunogen**

A synthetic phosphorylated peptide around T185/Y187 of human ERK2 (NP\_002736.3).

#### **Synonyms**

### **Contact**

•

www.abclonal.com

## **Product Information**

**Source** Rabbit Isotype

**Purification** 


IgG Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of lysates from Jurkat cells, using Phospho-ERK1-T202/Y204 + ERK2-T185/Y187 pAb (AP1120) at 1:1000 dilution or ERK1 / ERK2 antibody (A16686). Jurkat cells were treated by PMA/TPA (200 nM) at  $37^{\circ}$ C for 10 minutes.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% BSA.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.