Leader in Biomolecular Solutions for Life Science

# Phospho-AMPKa1/AMPKa2-T183/T172 Rabbit pAb



Catalog No.: AP1171

3 Publications

## **Basic Information**

# **Observed MW**

62kDa

#### **Calculated MW**

64kDa/65kDa/62kDa

### Category

Mouse Monoclonal Antibody

#### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

The protein encoded by this gene belongs to the ser/thr protein kinase family. It is the catalytic subunit of the 5'-prime-AMP-activated protein kinase (AMPK). AMPK is a cellular energy sensor conserved in all eukaryotic cells. The kinase activity of AMPK is activated by the stimuli that increase the cellular AMP/ATP ratio. AMPK regulates the activities of a number of key metabolic enzymes through phosphorylation. It protects cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. Alternatively spliced transcript variants encoding distinct isoforms have been observed.

# **Recommended Dilutions**

WB

1:500 - 1:2000

# Immunogen Information

**Gene ID** 5562/5563

**Swiss Prot** Q13131/P54646

#### **Immunogen**

A synthetic phosphorylated peptide around T183 of human AMPKa1/AMPKa2PRKAA1 (NP 006242.5).

## **Synonyms**

AMPKa1/AMPKa2; Phospho-AMPKa1/AMPKa2-T183/T172

## **Contact**

•

www.abclonal.com

# **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of various lysates using Phospho-AMPKa1/AMPKa2-T183/T172 Rabbit pAb (AP1171) at 1:1000 dilution.293T cells were treated by oligomycin (0.5 uM) at 37°C for 30 minutes.C2C12 cells were treated by oligomycin (0.5 uM) at 37°C for 30 minutes.C6 cells were treated by oligomycin (0.5 uM) at 37°C for 30 minutes after serum-starvation overnight.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% BSA.

Detection: ECL Enhanced Kit (RM00021).

Exposure time: 60s.