AP1439

Leader in Biomolecular Solutions for Life Science

# Phospho-Myosin Light Chain 2-T18 Rabbit mAb

ABclonal<sup>®</sup>

Catalog No.: AP1439 Recombinant

## **Basic Information**

Observed MW 18kDa

Calculated MW 20kDa

**Category** SMab Recombinant Monoclonal Antibody

Applications WB,ELISA

Cross-Reactivity Human

CloneNo number ARC62064

## Background

Myosin, a structural component of muscle, consists of two heavy chains and four light chains. The protein encoded by this gene is a myosin light chain that may regulate muscle contraction by modulating the ATPase activity of myosin heads. The encoded protein binds calcium and is activated by myosin light chain kinase. Two transcript variants encoding different isoforms have been found for this gene.

## **Recommended Dilutions**

### Immunogen Information

WB

1:500 - 1:1000

**Gene ID** 10398

Swiss Prot P24844

#### Immunogen

A synthetic phosphorylated peptide around T18 of human Myosin Light Chain 2 []NP\_006088.2[].

#### Synonyms

LC20; MLC2; MRLC1; MYRL2; MLC-2C; MMIHS4; Phospho-Myosin Light Chain 2-T18

## Contact

# **Product Information**

www.abclonal.com

**lsotype** IgG Purification Affinity purification


#### Storage

Source

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of HeLa, using Phospho-Myosin Light Chain 2-T18 Rabbit mAb (AP1439) at 1:1000 dilution.HeLa cells were treated by Calyculin A (100 nM) at 37°C for 30 minutes after serum-starvation overnight. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 90s.