Leader in Biomolecular Solutions for Life Science



HRP-conjugated Goat anti-Rabbit IgG Heavy Chain

Catalog No.: AS063 42 Publications

Basic Information

Observed MW 55kDa

Calculated MW

Category Secondary Antibody

Applications

WB

Cross-Reactivity

Conjugate

HRP

Background

Secondary antibodies are affinity-purified antibodies which will work with target-specific primary antibody in the detection, sorting or purification of its specified target. Secondary antibodies offer increased versatility enabling users to use many detection systems (e.g. HRP, AP, fluorescence). They can also provide greater sensitivity through signal amplification as multiple secondary antibodies . Most commonly, secondary antibodies are generated by immunizing the host animal (different from host species of primary antibody) with a pooled population of normal immunoglobulins from the host species of primary antibody and can be further purified and modified (i.e. antibody fragmentation, label conjugation, etc.) to ensure well-characterized specificity to corresponding normal immunoglobulins.

Recommended Dilutions

WB 1:1000-10000

Immunogen Information

Gene ID Swiss Prot

Immunogen Rabbit IgG

Synonyms

Contact

•

www.abclonal.com

Product Information


Source Isotype
Goat IgG

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 50% glycerol,pH7.3.

PurificationAffinity purification

Validation Data



Western blot analysis of lysates from Rabbit and Mouse IgG, using HRP-conjugated Goat Anti-Rabbit IgG

Heavy Chain (AS063) at 1:5000 dilution. Lysates/proteins: 100ng per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.