

Recombinant Human IL-1 beta Protein

Catalog No.: RP00002 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	3553	P01584

Tags

C-His

Synonyms

IL-1; IL1-BETA; IL1F2; IL1 beta; IL1B

Product Information

Source	Purification
<i>E. coli</i>	> 95% by SDS-PAGE.

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

Background

Interleukin-1 beta (IL1 beta or IL1B) also known as catabolin, is a member of the interleukin 1 cytokine family. This cytokine is produced by activated macrophages as a proprotein, which is proteolytically processed to its active form by caspase 1 (CASP1/ICE). This cytokine is an important mediator of the inflammatory response, and is involved in a variety of cellular activities, including cell proliferation, differentiation, and apoptosis. The induction of cyclooxygenase-2 (PTGS2/COX2) by this cytokine in the central nervous system (CNS) is found to contribute to inflammatory pain hypersensitivity.

Basic Information

Description

Recombinant Human IL-1 beta Protein is produced by *E. coli* expression system. The target protein is expressed with sequence (Ala117-Ser269) of human IL-1 beta (Accession #NP_000567.1) fused with an initial Met at the N-terminus and a 6xHis tag at the C-terminus.

Bio-Activity


1. Recombinant Human IL1β protein was measured by NF-κB reporter assay in HEK293 cells. The ED₅₀ for this effect is 1-10 pg/mL, corresponding to a specific activity of 1×10⁸~1×10⁹ units/mg. 2. Measured in a cell proliferation assay using D10.G4.1 mouse helper T cells. The ED₅₀ for this effect is 0.217-0.870 ng/mL, corresponding to a specific activity of 1.15×10⁶~4.61×10⁶ units/mg. 3. Measured by its binding ability in a functional ELISA. Immobilized Human IL-1 beta (Catalog: RP00002) at 10 μg/mL (100 μL/well) can bind Biotinylated Human IL-1R1 (Catalog: RP01036) with a linear range of 0.002-1 μg/mL.

Storage


Store the lyophilized protein at -20°C to -80 °C for long term. After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week. Avoid repeated freeze/thaw cycles.

Contact


Validation Data




Recombinant Human IL-1 beta Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 18 kDa.



Recombinant Human IL1 β protein was measured by NF- κ B reporter assay in HEK293 cells. The ED₅₀ for this effect is 1-10 pg/mL, corresponding to a specific activity of $1 \times 10^8 \sim 1 \times 10^9$ units/mg.



Recombinant Human IL-1 beta promotes the proliferation of D10.G4.1 mouse helper T cells. The ED₅₀ for this effect is 0.217-0.870 ng/mL, corresponding to a specific activity of $1.15 \times 10^6 \sim 4.60 \times 10^6$ units/mg.



Immobilized Human IL-1 beta (Catalog: RP00002) at 10 μ g/mL (100 μ L/well) can bind Biotinylated Human IL-1R1 (Catalog: RP01036) with a linear range of 0.002-1 μ g/mL.