Leader in Biomolecular Solutions for Life Science

Recombinant Human EphA2/ECK Protein

Catalog No.: RP00024 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 1969 P29317

numai

Tags No tag

Synonyms

ARCC2;CTPA;CTPP1;CTRCT6;ECK;EPHA2

Product Information

Source Purification

Baculovirus-Infected > 95% by SDS-Sf9 Cells PAGE.

Endotoxin

< 1.0 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

Ð

www.abclonal.com

Background

Eph receptor A2 (Ephrin type-A receptor 2 or EphA2) is a member of the ephrin receptor subfamily of the protein-tyrosine kinase family.The receptor tyrosine kinase which binds promiscuously membrane-bound ephrin-A family ligands residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Activated by the ligand ephrin-A1/EFNA1 regulates migration, integrin-mediated adhesion, proliferation and differentiation of cells. Regulates cell adhesion and differentiation through DSG1/desmoglein-1 and inhibition of the ERK1/ERK2 (MAPK3/MAPK1, respectively) signaling pathway. May also participate in UV radiation-induced apoptosis and have a ligand-independent stimulatory effect on chemotactic cell migration.

Basic Information

Description

Recombinant Human EphA2/ECK Protein is produced by insect cell-baculovirus expression system. The target protein is expressed with sequence (Asp596-Ile976) of human EphA2 (Accession #NP_004422.2).

Bio-Activity


Measured by its binding ability in a functional ELISA. Immobilized Human EphA2 at 1 μ g/mL (100 μ L/well) can bind Human EFNA1 with a linear range of 4.883 ng/mL-9.206 μ g/mL.


Storage

Store the lyophilized protein at -20°C to -80 °C for long term.
br>After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.
Avoid repeated freeze/thaw cycles.



Validation Data





Immobilized Human EphA2 at 1 $\mu g/mL$ (100 $\mu L/well) can bind Human EFNA1 with a linear range of 4.883 ng/mL-9.206 <math display="inline">\mu g/mL.$

Recombinant Human EphA2/ECK Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 43 kDa.