RP00050

Leader in Biomolecular Solutions for Life Science

Recombinant Human IL-11 Protein

Catalog No.: RP00050 Recombinant

Swiss Prot

P20809



Sequence Information

Gene ID

3589

Species Human

Tags No tag

Synonyms

IL11;AGIF;IL-11

Product Information

Source <I>E. coli</I> Purification > 97% by SDS-PAGE.

Endotoxin

< 1.0 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of 20mM Tris, 150mM NaCl, pH 8.0.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

€

www.abclonal.com

Background

The protein is a member of the gp130 family of cytokines. These cytokines drive the assembly of multisubunit receptor complexes, all of which contain at least one molecule of the transmembrane signaling receptor IL6ST (gp130). This cytokine is shown to stimulate the T-cell-dependent development of immunoglobulin-producing B cells. It is also found to support the proliferation of hematopoietic stem cells and megakaryocyte progenitor cells.

Basic Information

Description

Recombinant Human IL-11 Protein is produced by <I>E. coli</I> expression system. The target protein is expressed with sequence (Pro22-Leu199) of human IL-11 (Accession #NP_000632.1) fused with an initial Met at the N-terminus.


Bio-Activity

Measured in a cell proliferation assay using TF-1 human erythroleukemic cells. The ED₅₀ for this effect is 1.3-5 ng/mL, corresponding to a specific activity of 2×10 ⁵-7.69×10⁵-units/mg.

Storage

Store the lyophilized protein at -20°C to -80 °C for long term.
br>After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.
Avoid repeated freeze/thaw cycles.

Validation Data



0.50 0.45 0.40 0.00

Recombinant human IL11 promotes the proliferation of TF-1 human erythroleukemic cells. The ED₅₀ for this effect is 1.3-5ng/mL, corresponding to a specific activity of 2×10^5 -7.69 $\times 10^5$ units/mg.

Active Recombinant Human IL-11 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 21 kDa.