

RP00113

Leader in Biomolecular Solutions for Life Science



Recombinant Human IgA Fc receptor/CD89 Protein

Catalog No.: RP00113 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	2204	P24071

Tags

C-hFc&His

Synonyms

FCAR;CD89;CTB-61M7.2;FcalphaRI

Background

This protein is a member of the immunoglobulin gene superfamily and receptor for the Fc region of IgA. The receptor is a transmembrane glycoprotein present on the surface of myeloid lineage cells such as neutrophils, monocytes, macrophages, and eosinophils, where it mediates immunologic responses to pathogens. It interacts with IgA-opsonized targets and triggers several immunologic defense processes, including phagocytosis, antibody-dependent cell-mediated cytotoxicity, and stimulation of the release of inflammatory mediators. Multiple alternatively spliced transcript variants encoding different isoforms have been described for this gene.

Basic Information

Description

Recombinant Human IgA Fc receptor/CD89 Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Gln22-Asn227) of human CD89/FCAR (Accession #NP_001991.1) fused with an Fc, 6xHis tag at the C-terminus.

Bio-Activity

Storage

Store the lyophilized protein at -20°C to -80 °C for long term. After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week. Avoid repeated freeze/thaw cycles.

Product Information

Source	Purification
HEK293 cells	> 97% by SDS-PAGE.

Endotoxin

< 0.1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution


Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact



www.abclonal.com

Validation Data



Recombinant Human IgA Fc receptor/CD89 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 65-75 kDa.