RP00118

Leader in Biomolecular Solutions for Life Science

Recombinant Human MIC-B Protein

Catalog No.: RP00118 Recombinant

Sequence Information

Background

SpeciesGene IDSwiss ProtHuman4277Q29980

Tags C-hFc&His

Synonyms

MICB;PERB11.2

Product Information

Source Purification HEK293 cells > 97% by SDS-PAGE.

Endotoxin

< 0.1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μ m filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

Ð

www.abclonal.com

This protein is a heavily glycosylated protein which is a ligand for the NKG2D type II receptor. Binding of the ligand activates the cytolytic response of natural killer (NK) cells, CD8 alphabeta T cells, and gammadelta T cells which express the receptor. This protein is stress-induced and is similar to MHC class I molecules; however, it does not associate with beta-2-microglobulin or bind peptides. Alternative splicing results in multiple transcript variants.

Basic Information

Description

Recombinant Human MIC-B Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Ala23-Gly298) of human MICB (Accession $\#NP_005922.2$) fused with an Fc, 6×His tag at the C-terminus.


Bio-Activity


Storage

Store the lyophilized protein at -20°C to -80 °C for long term.
br>After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week. Avoid repeated freeze/thaw cycles.



Validation Data





The purity of Human MICB Protein (Cat.RP00118) was greater than 95% as determined by SEC-HPLC.

Recombinant Human MIC-B Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at approximately 75 kDa.