Leader in Biomolecular Solutions for Life Science

Recombinant Human Ephrin-A4/EFNA4 Protein

Catalog No.: RP00143 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 1945 P52798

Human

Tags C-hFc&His

-

Synonyms EFL4;EPLG4;LERK4;EFNA4

Product Information

Source HEK293 cells

Purification > 80% by SDS-PAGE.

Endotoxin

< 0.1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

Ð

www.abclonal.com

Background

EPH-related receptor tyrosine kinase ligand 4 (Ephrin-A4) also known as EFNA4, is a member of the Ephrin family. The Eph family receptor interacting proteins (ephrins) are a family of proteins that serve as the ligands of the Eph receptor, which compose the largest known subfamily of receptor protein-tyrosine kinases (RTKs). Eph/ephrin interactions are implicated in axon guidance, neural crest cell migration, establishment of segmental boundaries, and formation of angiogenic capillary plexi. Ephrin subclasses are further distinguished by their mode of attachment to the plasma membrane: ephrin-A ligands bind EphA receptors and are anchored to the plasma membrane via a glycosylphosphatidylinositol (GPI) linkage, whereas ephrin-A ligands bind EphB receptors and are anchored via a transmembrane domain. Ephrin-A4/EFNA4 functions as a cell surface GPI-bound ligand for Eph receptor, a family of receptor tyrosine kinases which are crucial for migration, repulsion and adhesion during neuronal, vascular and epithelial development.

Basic Information

Description

Recombinant Human Ephrin-A4/EFNA4 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Leu26-Gly171) of human Ephrin-A4/EFNA4 (Accession # NP_005218.1) fused with an Fc, 6×His tag at the C-terminus.

Bio-Activity


Measured by its binding ability in a functional ELISA. Immobilized Human EphA7 at 0.5 μ g/mL (100 μ L/well) can bind Human EFNA4 with a linear range of 0.039-0.942 ng/mL.


Storage

Store the lyophilized protein at -20°C to -80 °C for long term.
 After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week. Avoid repeated freeze/thaw cycles.



Validation Data





Immobilized Human EphA7 at 0.5 $\mu g/mL$ (100 $\mu L/well)$ can bind Human EFNA4 with a linear range of 0.039-0.942 ng/mL.

Active Recombinant Human Ephrin-A4/EFNA4 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 50 kDa.