Leader in Biomolecular Solutions for Life Science

Recombinant Human Osteoprotegerin/TNFRSF11B/OPG Protein



Catalog No.: RP00459 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 4982 000300

Tags C-6×His

_

Synonyms

TNFRSF11B; OCIF; OPG; PDB5; TR1; TNF receptor superfamily member 11b;Osteoprotegerin;OCIF;OPG;PDB5;TR 1

Product Information

Source Purification
HEK293 cells > 95% by SDSPAGE.

Endotoxin

< 1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.2 µm filtered solution of 20 mM PB, 150 mM NaCl, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water.

Background

This protein is a member of the TNF-receptor superfamily. This protein is an osteoblast-secreted decoy receptor that functions as a negative regulator of bone resorption. This protein specifically binds to its ligand, osteoprotegerin ligand, both of which are key extracellular regulators of osteoclast development. Studies of the mouse counterpart also suggest that this protein and its ligand play a role in lymphnode organogenesis and vascular calcification. Alternatively spliced transcript variants of this gene have been reported, but their full length nature has not been determined.

Basic Information

Description

Recombinant Human Osteoprotegerin/TNFRSF11B/OPG Protein is produced by Human Cell expression system. The target protein is expressed with sequence (Glu22-Leu401) of human Osteoprotegerin/TNFRSF11B/OPG (Accession #000300) fused with a 6×His tag at the C-terminus.

Bio-Activity

Storage


Store the lyophilized protein at -20°C to -80 °C for long term.
br>After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.
Avoid repeated freeze/thaw cycles.

Contact



www.abclonal.com

Validation Data



Recombinant protein Human Osteoprotegerin/TNFRSF11B/OPG was determined by SDS-PAGE under reducing conditions with Coomassie Blue, showing a band at 60 kDa.