Leader in Biomolecular Solutions for Life Science

# ABclonal www.abclonal.com

## Recombinant Human PD-1/PDCD1/CD279 Protein

Catalog No.: RP00552 Recombinant

## **Sequence Information**

**Species Gene ID Swiss Prot** Human 5133 Q15116

## Tags

C-mFc

#### **Synonyms**

PDCD1; CD279; PD-1; PD1; SLEB2; hPD-1; hPD-I; hSLE1; programmed cell death 1;PD-1;CD279;PD1;SLEB2;hPD-1;hPD-I;hSLE1;PD-1/CD279

#### **Product Information**

**Source** Purification HEK293 cells > 95% by SDS-

PAGE.

#### **Endotoxin**

< 1 EU/µg of the protein by LAL method.

#### **Formulation**

Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.2.Contact us for customized product form or formulation.

#### Reconstitution

Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water.

### **Background**

Programmed cell death protein 1(PDCD1) is a single-pass type I membrane protein and contains 1 Ig-like V-type domain. PD-1 is a member of the extended CD28/CTLA-4 family of T cell regulators. PDCD1 inhibits the T-cell proliferation and production of related cytokines including IL-1, IL-4, IL-10 and IFN-  $\gamma$  by suppressing theactivation and transduction of PI3K/AKT pathway. In addition, coligation of PDCD1 inhibits BCR-mediatingsignal by dephosphorylating key signal transducer. PDCD1 has been suggested to be involved in lymphocyteclonal selection and peripheral tolerance, and thus contributes to the prevention of autoimmune diseases. As acell surface molecule, PDCD1 regulates the adaptive immune response. Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function.

#### **Basic Information**

#### **Description**

Recombinant Human PD-1/PDCD1/CD279 Protein is produced by Human cells expression system. The target protein is expressed with sequence (Leu25-Gln167) of human PD-1/PDCD1/CD279 (Accession #Q15116) fused with an mFc tag at the C-terminus.

#### **Bio-Activity**

#### Storage


Store the lyophilized protein at -20  $^{\circ}$ C to -80  $^{\circ}$ C for long term.<br/>
hr>After reconstitution, the protein solution is stable at -20  $^{\circ}$ C for 3 months, at 2-8  $^{\circ}$ C for up to 1 week. Avoid repeated freeze/thaw cycles.

#### **Contact**



www.abclonal.com

## **Validation Data**



Recombinant Human PD-1/PDCD1/CD279 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.