Leader in Biomolecular Solutions for Life Science



Recombinant Mouse Fc-gamma RI/CD64 Protein

Catalog No.: RP00710 Recombinant

Sequence Information

Species Gene ID Swiss Prot Mouse 14129 P26151

Tags

C-His

Synonyms

High affinity immunoglobulin gamma Fc receptor I; IgG Fc receptor I; Fc-gamma RI; FcRI; CD64; Fcgr1

Product Information

Source Purification HEK293 cells > 95% by SDS-PAGE.

Endotoxin

< 1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact



www.abclonal.com

Background

CD64, also known as Fc-gamma receptor 1 (Fc γ RI), is a type of integral membrane glycoprotein that bindsmonomeric IgG-type antibodies with high affinity. After binding IgG, CD64 interacts with an accessory chainknown as the common γ chain (γ chain), which possesses an ITAM motif that is necessary for triggering cellularactivation. CD64 is composed of a signal peptide, three extracellular immunoglobulin domains of the C2-typeused to bind antibody, a hydrophobic transmembrane domain, and a short cytoplasmic tail. CD64 mediatesendocytosis, phagocytosis, antibody-dependent cellular cytotoxicity, cytokine release, and superoxideproduction. It is normally expressed on the surfaces of monocytes and macrophages.

Basic Information

Description


Recombinant Mouse Fc γ RI/FCGR1A/CD64 Protein is produced by Human Cells expression system. The target protein is expressed with sequence (Glu25-Pro297) of mouse Fc γ RI/FCGR1A/CD64 (Accession #P26151) fused with a 6×His tag at the C-terminus.

Bio-Activity

Storage

Store the lyophilized protein at -20°C to -80 °C for long term.
br>After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.
Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Mouse Fc-gamma RI/CD64 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.