Leader in Biomolecular Solutions for Life Science

# **Recombinant Human CD44 Protein**

Catalog No.: RP00973 Recombinant 2 Publications



## **Sequence Information**

**Species Gene ID Swiss Prot** Human 960 P16070-1

# Tags

C-hFc&His

#### **Synonyms**

CD44; CDW44; CSPG8; ECMR-III; HCELL; HUTCH-I; IN; LHR; MC56; MDU2; MDU3; MIC4; Pgp1; CD44 antigen; CDW44; CSPG8; ECMR-III; HCELL; HUTCH-I; IN; LHR; MC56; MDU2; MDU3; MIC4; Pgp1; C D44

## **Product Information**

Source Purification
HEK293 cells > 97% by SDSPAGE.

### **Endotoxin**

 $< 1.0 \; \text{EU/}\mu\text{g}$  of the protein by LAL method.

#### **Formulation**

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

# Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## **Contact**



www.abclonal.com

## **Background**

This protein is a cell-surface glycoprotein involved in cell-cell interactions, cell adhesion and migration. It is a receptor for hyaluronic acid (HA) and can also interact with other ligands, such as osteopontin, collagens, and matrix metalloproteinases (MMPs). This protein participates in a wide variety of cellular functions including lymphocyte activation, recirculation and homing, hematopoiesis, and tumor metastasis. Transcripts for this gene undergo complex alternative splicing that results in many functionally distinct isoforms, however, the full length nature of some of these variants has not been determined. Alternative splicing is the basis for the structural and functional diversity of this protein, and may be related to tumor metastasis.

### **Basic Information**

#### Description

Recombinant Human CD44 Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Gln21-Pro220) of human CD44 (Accession #NP 000601.3) fused with an Fc, 6×His tag at the C-terminus.


#### **Bio-Activity**

Measured by its binding ability in a functional ELISA. Immobilized Hyaluronic Acid at  $100 \mu g/mL$  ( $100 \mu L/well$ ) can bind Human CD44 with a linear range of 20-330 ng/mL.


#### Storage

Store the lyophilized protein at -20°C to -80 °C for long term. <br/>br>After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.<br/>Avoid repeated freeze/thaw cycles.

# **Validation Data**



Recombinant Human CD44 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 70-85 kDa.



Immobilized Hyaluronic Acid at  $100 \mu g/mL$  ( $100 \mu L/well$ ) can bind Recombinant Human CD44 with a linear range of 20-330 ng/mL.