

RP00989

Leader in Biomolecular Solutions for Life Science



Recombinant Human Osteopontin/SPP-1 Protein

Catalog No.: RP00989

Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
HEK293 cells	6696	P10451-5

Tags

C-His

Synonyms

BNSP;BSPI;ETA-1;OPN;Osteopontin;SPP1

Product Information

Source	Purification
	> 87% by SDS-PAGE.

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

Background

Basic Information

Description

Recombinant Human Osteopontin/SPP-1 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ile17-Asn300) of human Osteopontin (Accession #NP_000573.1) fused with a 6×His tag at the C-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized Human SPP1 Protein at 1 μg/mL (100 μL/well) can bind Osteopontin Rabbit mAb with a linear range of 0.976-11.58 ng/mL.

Storage


Store the lyophilized protein at -20°C to -80 °C for long term. After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week. Avoid repeated freeze/thaw cycles.

Contact




www.abclonal.com


Validation Data



Recombinant Human Osteopontin/SPP-1 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 55 kDa.



Immobilized recombinant Human SPP1 Protein at 1µg/mL (100 µL/well) can bind Osteopontin Rabbit mAb with a linear range of 0.976-11.58ng/mL.



Recombinant Human Osteopontin/SPP-1 immobilized protein to support the adhesion of B16 cells. When 5×10^4 cells/well are added to a Recombinant Human (rh) Osteopontin/OPN coated plate, cell adhesion is enhanced in a dose-dependent manner after 1 hour incubation at 37 °C. The ED_{50} for this effect is 23.9-95.7 ng/mL.